

Psych CE

An Introduction to Severe Psychopathology

1. People with schizophrenia

- A. are more dangerous than those without a psychotic disorder
 - B. have multiple personalities
 - C. are frequently homeless or are hospitalized
 - D. live with their families, in supportive housing, or independently.
-

2. Negative symptoms include

- A. loss of motivation & difficulty showing emotions
 - B. delusions & hallucinations
 - C. difficulty with memory & learning
 - D. illogical thinking & body tics
-

3. Treatment of psychotic disorders may include all of the following except

- A. antipsychotic medication
 - B. psychosocial treatment
 - C. genetic testing
 - D. substance use treatment
-

4. The psychotic disorder that is a combination of schizophrenia and mood disorder is

- A. Schizophreniform Disorder
 - B. Schizoaffective Disorder
 - C. Delusional Disorder
 - D. Unspecified Schizophrenia Spectrum Disorder
-

5. Schizophreniform is different from schizophrenia in that

- A. it last for less than 6 months
 - B. the person also has bipolar symptoms
 - C. there are no hallucinations
 - D. the person can distinguish between what is real and imagined
-

6. With Brief Psychotic Disorder the person will return to normal functioning within

- A. a year

- B. six months
 - C. three months
 - D. one month
-

7. Believing someone is in love with them to the point they will not stop attempts at contacting the person is this type of delusion

- A. Jealous
 - B. Grandios
 - C. Erotomaniac
 - D. Persecutory
-

8. Episodes of this type of psychosis are common in emergency departments

- A. Schizophrenia
 - B. Substance-induced Psychotic Disorder
 - C. Delusional Disorder
 - D. Brief Psychotic Disorder
-

9. Potential causes of psychosis include all except

- A. Genetics
 - B. Environmental Factors
 - C. Neurotransmitters
 - D. Education
-

10. One of the most important pieces for successful recovering or living with a psychotic disorder is

- A. early treatment at first onset
 - B. understanding family predisposition
 - C. electroconvulsive therapy
 - D. placement in group home
-