

# Psych CE

## Depression in New Mothers

1. The baby blues are experienced by what percentage of women?

- A. 50-60 %
  - B. 60-70%
  - C. 70-80%
  - D. 80-90%
- 

2. According to the author, risk factors for postpartum depression may include each of the following EXCEPT:

- A. Experiencing postpartum depression in a previous pregnancy, a family history of depression or mood disorders or a stressful life
  - B. Pregnancy complications, a baby born with special needs or health, or twins, triplets, or multiples born
  - C. A diagnosis of an anxiety disorder, a personality disorder, or a co-occurring mental health/substance abuse issue
  - D. Having issues in your relationship, having a compromised immune system, or experiencing financial hardship
- 

3. The baby blues is equally as dangerous as postpartum depression for most women.

- A. True
  - B. False
- 

4. Which attachment style do people strive for with their children?

- A. Secure
  - B. Avoidant
  - C. Anxious
  - D. Disorganized
- 

5. Postpartum psychosis tends to be much more dangerous than postpartum depression.

- A. True
  - B. False
-

**6. A multidisciplinary treatment team involves which of the following providers?**

- A. Therapist**
  - B. Physician**
  - C. Psychiatrist**
  - D. All of the above**
- 

**7. Women with depression are more likely to misuse substances?**

- A. True**
  - B. False**
- 

**8. According to the text, the following risk factors pose a higher chance of experiencing baby blues EXCEPT for:**

- A. A family history of depression**
  - B. Bipolar disorder**
  - C. Addiction to alcohol**
  - D. Pregnancy complications**
- 

**9. Interpersonal therapy, which has been found to be clinically beneficial for women with postpartum depression, explores four different interpersonal problems, including role transition, role dispute, interpersonal deficits, and:**

- A. Grief**
  - B. Unwanted thoughts and feelings**
  - C. Lack of personal acceptance**
  - D. Relationship issues**
- 

**10. Of the following treatment services, what is most likely one that should be started right away for women with postpartum depression?**

- A. Medication**
  - B. Group therapy**
  - C. Family therapy**
  - D. Individual therapy**
- 

**11. What should be completed as soon as possible once a provider begins working with a new mother with depression?**

- A. A long term plan for what recovery looks like**
- B. Safety plan**
- C. Goal setting**

**D. Cope ahead plan**

---

**12. The diagnosis of postpartum depression will involve a screening tool for depression.**

- A. True**
  - B. False**
- 

**Copyright © 2020 Psych CE**

**Visit us at <https://www.psychce.com>**